

GERM CELL DIFFERENTIATION AND SEXUAL MATURATION OF THE
FEMALE *NEPTUNEA (BARBITONIA) ARTHRITICA CUMINGII* (CROSSE, 1862)
(GASTROPODA: BUCCINIDAE)

Ee-Yung Chung¹, Sung Yeon Kim², Gab-Man Park³ & Jong Man Yoon⁴

ABSTRACT

Oogenesis, the gonadosomatic index (GSI), reproductive cycle, and first sexual maturation of the female *Neptunea (Barbitonia) arthritica cumingii* have been investigated by light and electron microscope observations. In the early vitellogenic oocyte, the Golgi complex and mitochondria were involved in the formation of glycogen, lipid droplets, and yolk granules. In late vitellogenic oocytes, the rough endoplasmic reticulum and multivesicular bodies were involved in the formation of proteid yolk granules in the cytoplasm. In particular, compared with the results of other gastropods, it differs in that appearances of cortical granules at the cortical layer and microvilli on the vitelline envelope, which is associated with heterosynthetic vitellogenesis, were not observed in vitellogenic oocytes during oogenesis. A mature yolk granule was composed of three components: main body (central core), superficial layer, and the limiting membrane. Monthly changes in the gonadosomatic index in females studied in 2002 and 2003 were closely associated with ovarian developmental phases. Spawning occurred between May and August in 2002 and 2003, and the main spawning occurred between June and July, when the seawater temperature rose to approximately 18–23°C. The female reproductive cycle can be classified into five successive stages: early active stage (September to October), late active stage (November to February), ripe stage (February to June), partially spawned stage (May to August), and recovery stage (June to August). The rate of individuals reaching the first sexual maturity was 53.1% in females of 51.0 to 60.9 mm in shell height, and 100% in those over 61.0 mm.

Key words: *Neptunea (Barbitonia) arthritica cumingii*, oogenesis, germ cell differentiation, sexual maturation.

INTRODUCTION

Neptunea arthritica cumingii (Crosse, 1862) is one of the most important edible gastropods in such East Asian countries as Korea, Japan, China, and Russia (Yoo, 1976; Kwon et al., 1993). This species is especially found in silty sand of the subtidal zone of the west coast of Korea. Recently, as the standing stock of this species gradually decreased due to extensive reclamation projects and reckless overharvesting, it has been designated as one of the important organisms in need of resource management.

There have been some studies on *Neptunea* species on aspects of reproduction, including the reproductive cycle (Takahashi et al., 1972; Takamaru & Fuji, 1981; Fujinaga, 1985; Kawai et al., 1994), and spawning (Miyawaki, 1953;

Amio, 1963; Son, 2003), on aspects of ecology including distribution (Ito & Tachizawa, 1981; Ito, 1982; Kwon et al., 1993), growth (MacIntosh & Paul, 1977; Fujinaga, 1987; Suzuki et al., 1996) of *N. arthritica*, and feeding (Pearce & Thorson, 1967) of *N. antiqua*. There has been one study on the spawning season of *N. cumingii* in the East China Sea (Amio, 1963). But, there are still gaps in our knowledge of its reproductive biology. So far, little information is available on ultrastructural study on germ cell differentiation and sexual maturation of *N. arthritica cumingii* in the Korean waters and the Japan Sea (Chung & Kim, 1996). However, there is some information on ultrastructural study of oogenesis in other gastropods (McCann-Coillier, 1977, 1979; Griffond & Gomot, 1979; Griffond, 1980; Hodgson & Eckelbarger, 2000; Pal & Hodgson, 2002).

¹School of Marine Life Science, Kunsan National University, Kunsan 573-701, Korea; eychung@kunsan.ac.kr

²National Fisheries Development Institute, Busan 690-902, Korea

³Department of Parasitology, Kwandong University College of Medicine, Gangnung 210-710, Korea

⁴Department of Aquatic Life Medicine, Kunsan National University, Kunsan 573-701, Korea

Therefore, the results of ultrastructural studies on germ cell differentiation of this species and other gastropods provides important information on the reproductive mechanisms. The reproductive cycles of the local populations in marine gastropods vary with such environmental factors as water temperature and food availability (Chung et al., 2002). Understanding the reproductive cycle and the spawning period of *N. arthritica cumingii* will provide necessary information for natural spat collections or the recruitment period and age determination of this population. In addition, data on first sexual maturity and reproductive strategy of this population are very useful information for natural resource management. The main aim of the present study is to understand germ cell differentiation during oogenesis, the reproductive cycle and first sexual maturity of this species.

MATERIALS AND METHODS

Sampling

Specimens of *Neptunea arthritica cumingii* (Crosse, 1862) were collected monthly at the subtidal zone of Maldo, Kunsan, Korea, from

January to December 2002 (Fig. 1). Snails ranging from 41.0 to 106.8 mm in shell height were used for the present study. After the snails were transported alive to the laboratory, shell heights were immediately measured.

Gonadosomatic Index (GSI)

A total of 486 individuals were used for calculation of the GSI. Monthly changes in the mean gonadosomatic index (GSI) were calculated by the following equation (Chung et al., 2002) (Fig. 2):

$$\text{GSI} = \frac{\text{Thickness of the gonad} \times 100}{\text{Diameter of posterior appendage including the gonad and digestive gland}}$$

Germ Cell Differentiation by Electron Microscopic Observation

For electron microscopical observations, excised pieces of the gonads were cut into small pieces and immediately fixed in 2.5% paraformaldehyde-glutaraldehyde in 0.1 M phosphate buffer (pH 7.4) for 2 h at 4°C. After initial fixation, the specimens were washed several times with the same buffer and then




FIG. 1. Map showing the sampling area.




FIG. 2. Anatomy of *Neptunea arthritica cumingii*, removed from its shell. Posterior appendage showing the ovary and digestive gland. X, Y and Z denote the sections for measurement of GSI. Three sections are spaced equally. Abbreviations: DG, digestive gland; OV, ovary; ST, stomach; SC, stomachal caecum.

further fixed in 1% osmium tetroxide dissolved in 0.2 M phosphate buffer solution (pH 7.4) for 1 h at 4°C. Specimens were then dehydrated in a series of increasing concentrations of ethanol, cleared in propylene oxide, and embedded in Epon-Araldite mixture. Ultrathin sections of Epon-embedded specimens were cut with glass knives with a Sorvall MT-2 microtome and an LKB ultramicrotome at a thickness of about 800–1,000 Å. Tissue sections were mounted on collodion-coated copper grids, stained with uranyl acetate followed by lead citrate, and examined with a JEM 100 CX-2 (80 kV) electron microscope.

Gonadal Development by Histological Observations

For light microscopic examination of histological preparations, a total of 456 individuals were used for histological analysis of the gonads from January to December 2002. Gonad tissues were removed from shells and preserved in Bouin's fixative for 24 h and then washed with running tap water for 24 h. Tissues were then dehydrated in alcohol and embedded in paraffin molds. Embedded tissues were sectioned at 5–7 µm thickness using a rotary microtome. Sections were mounted on glass slides, stained with Hansen's hema-

toxylin-0.5% eosin, Mallory's triple stain and PAS stain, and examined using a light microscope.

First Sexual Maturity

The first sexual maturation of a total of 187 female individuals (31.4–90.5 mm in shell height) were investigated histologically in order to determine the shell heights of snails reaching maturation and participating in reproduction from May (ripe stage) to late August (after spawning).

RESULTS

Position and Morphology of the Gonads

Neptunea arthritica cumingii is a dioecious species composed of well-defined female and male individuals. The ovary is located on the surface of the digestive gland in the spiral posterior region of the shell (Fig. 2). The ovary is composed of numerous oogenic follicles. As the ovary matured, it extended to the outer part of the digestive gland. As maturation progresses, the sex of the snail can be distinguishable easily by color: the ovary being pale yellow and testis yellowish-brown. At this time,




FIG. 3. Monthly changes in the gonadosomatic index of female *Neptunea arthritica cumingii*, for two years from January 2002 to December 2003.

if it was slightly scratched with a razor, ripe eggs readily discharged from the ovary. But after spawning, the ovary degenerated, and it became difficult to distinguish their sexes by external color or dissection.

Monthly Changes in the Gonadosomatic Index (GSI)

Monthly GSI changes in females were showed in Figure 3. In 2002, the GSI slowly increased from September and reached the maximum (mean 3.11) in April when seawater temperature rapidly increased. The GSI rapidly decreased after May, and the values reached the minimum in August, when spawning was completely finished. Monthly changes in the GSI in 2003 showed similar patterns with those in 2002.

Germ Cell Differentiation in the Ovary by Electron Microscopic Observations

Ultrastructural observations allow the germ cell developmental phases during oogenesis can be divided into 4 phases: (1) oogonial phase, (2) previtellogenic phase, (3) vitellogenic phase, and (4) mature phase. Characteristic features in each stage were as follows:

Oogonial Phase: Oogonia in the oogonial phase, which propagated on the germinal epithelium (follicular wall), were oval and 15 μm in diameter. They commonly were single or formed a cluster on the germinal epithelium. Each oogonium had a large nucleus with chromatin, several mitochondria, and the endo-

plasmic reticulum, vacuoles in the cytoplasm (Fig. 4A).

Previtellogenic Phase: Previtellogenic oocytes were 25–90 μm in diameter. With cytoplasmic growth, several small mitochondria, a well-developed endoplasmic reticulum and several vacuoles were concentrated around the nucleus in the cytoplasm of the previtellogenic oocyte. The number of Golgi complexes, scattered from the perinuclear region to the cortical region of the oocyte, increased. At this time, many vacuoles formed by the Golgi complex appeared around the endoplasmic reticulum, several mitochondria, and large vesicles were present in the cytoplasm of the previtellogenic oocyte (Fig. 4B).

Vitellogenic Phase: In the early vitellogenic oocyte, especially, well-developed endoplasmic reticulum and vacuoles in the cytoplasm were concentrated around the nucleus having nucleoli. At this time, the follicle cell, which lied adjacent to the early vitellogenic oocyte, had an elongated nucleus. In particular, electron-dense granules and several lipid droplets were accumulated in the cytoplasm of the follicle cell (Fig. 4C). With the initiation of yolk formation, lipid droplets were accumulated in the vacuoles formed by the Golgi complex in the perinuclear region. Lipid droplets diffused toward the cortical layer, and then glycogen particles appeared around the mitochondria at the cortical region of early vitellogenic oocytes (Fig. 4D). At this time, after electron-dense materials were accumulated in the Golgi complex (Golgi sac, Golgi vacuoles, and Golgi vesicles), lipid droplets were formed by secre-




FIG. 4. Electron micrographs of the previtellogenic and early vitellogenic phases during oogenesis of *Neptunea arthritica cumingii* (A–F). A: Oogonia in the oogonial phase, with a large nucleus and several mitochondria in the cytoplasm; B: A previtellogenic oocyte, with a large nucleus with a few nucleolus and several mitochondria, the Golgi complex, and vacuoles in the cytoplasm; C: An early vitellogenic oocyte attached to a follicle cell, with a large nucleus containing chromatin and a number of vacuoles and well-developed endoplasmic reticulum in the cytoplasm; D: An early vitellogenic oocytes, with well-developed Golgi complex, glycogen particles and lipid droplets; E: An early vitellogenic oocyte, with lipid droplets formed by secretions in vacuoles and vesicles; F: An early vitellogenic oocyte, with a lipid droplet surrounded by the endoplasmic reticulum and the mitochondria. Abbreviations: CR, chromatin; ER, endoplasmic reticulum; G, Golgi complex; GS, Golgi sac; GV, Golgi vacuole; GVc, Golgi vesicle; LD, lipid droplet; M, mitochondrion; N, nucleus; NO, Nucleolus; NU, nucleolus; OC, oocyte; OG, oogonium; ER, Endoplasmic reticulum; Va, vacuole; Ve, vesicle.

tion of electron-dense materials in the large vacuoles and small vesicles, which were formed by the Golgi vacuoles and Golgi vesicles (Fig. 4E). On the other hand, relatively large lipid droplet was surrounded by the endoplasmic reticulum, the mitochondria and glycogen particles in the cytoplasm of the early vitellogenic oocyte (Fig. 4F). In the late vitellogenic oocyte, lots of yolk granules appeared between the rough endoplasmic reticulum and the mitochondria at the cortical layer in the cytoplasm (Fig. 5A). At this time, the multivesicular bodies, which were formed by the modified cristae of the mitochondria, ap-

peared near the nuclear envelope of the nucleus in the late vitellogenic oocyte. Yolk precursors, such as glycogen particles, lipid droplets, yolk granules and multivesicular bodies, were accumulated in the cytoplasm (Fig. 5B). Eventually, proteid yolk granules were formed by yolk granules and multivesicular bodies (Fig. 5C).

Mature Phase: Mature oocytes were about 180–250 × 300–450 μm in diameter. In the mature oocyte, various sizes of proteid yolk granules were intermingled with small lipid yolk granules, and it became a small mature yolk granule (Fig. 5C). Relatively small mature yolk




FIG. 5. Electron micrographs of late vitellogenic and mature phases during oogenesis of *Neptunea arthritica cumingii* (A–D). A: A late vitellogenic oocyte, with yolk granules between the rough endoplasmic reticulum and the the mitochondria; B: A late vitellogenic oocyte, with a number of multivesicular bodies formed by modified mitochondria; C: A late vitellogenic oocyte, with proteid yolk granules formed by yolk granules and multivesicular bodies; D: A mature oocyte, with a mature yolk granule being composed of the main body (central core), superficial layer and a limiting membrane of a yolk granule. Abbreviations: LD, lipid droplet; LM, limiting membrane; LYG, lipid yolk granule; M, mitochondrion; MB, main body; MM, modified mitochondrion; MYG, mature yolk granule; MV, multivesicular body; N, nucleus; NE, nuclear envelope; NU, nucleolus; PYG, proteid yolk granule; rER, rough endoplasmic reticulum; SL, superficial layer.




FIG. 6. Frequency of gonadal phases of *Neptunea arthritica cumingii* and the mean water temperatures, for two years, from January 2002 to December 2003.

granules were continuously mixed with each other and became large mature yolk granules in the cytoplasm. A fully mature yolk granule is composed of three components: (1) main body, (2) superficial layer, and (3) a limiting membrane (Fig. 5D).

Reproductive Cycle with the Gonad Developmental Stage

Based on the morphological features and sizes of germ cells and the tissue cells around them, the reproductive cycle with gonadal phases can be classified into five stages in females. Especially, the reproductive cycle and monthly changes in water temperatures showed similar patterns in 2002 and 2003 (Fig. 6). The criteria in defining of each stage are as follows:

Early Active Stage: The gonadal volume was small, and the follicles occupied approximately 25% of the gonad. The follicular walls were relatively thick. Oogonia and previtellogenic oocytes propagated along the oogenic follicular walls and mesenchymal tissues of the ovary. The oogonia and previtellogenic oocytes are

about 15–25 μm in size, respectively. At this time, early vitellogenic oocytes of 25–50 μm in diameter formed an egg-stalk attached to the walls (Fig. 7A). The individuals in the early active stage were found from September to October when seawater temperatures were gradually decreasing.

Late Active Stage: This stage is characterized by the presence of developing early vitellogenic oocytes. Follicular walls (germinal epithelium) were thin. A number of early vitellogenic oocytes of 100–140 μm in diameter were attached to the follicular walls through each egg-stalk. With the initiation of yolk formation, there were numerous yolk granules in the cytoplasm of late vitellogenic oocytes of 150–200 $\times$ 250–300 μm in diameter. Some fully mature oocytes were free in the lumen of the follicle (Fig. 7B, C). The individuals in the late active stage appeared from November to February.

Ripe Stage: In females, the majority of oocytes grew to 160–180 μm in diameter, occupied over 70% of the gonad, and follicular walls became very thin. Mature oocytes growing up to 180–250 $\times$ 300–450 μm in diameter became

tetragonal or polygonal in shape, and contained a number of mature yolk granules (Fig. 7D). Mature or ripe ovaries were found in February through June, when seawater temperatures gradually increased.

Partially Spawned Stage: Since about 50–70% of the oocytes in the follicles were discharged, the lumen of the follicles emptied. Spawned ovaries were characterized by the presence of a few undischarged vitellogenic




FIG. 7. Photomicrographs of the gonadal phases of female *Neptunea arthritica cumingii*. A: Transverse section of oogenic follicles in the early active stage; B: Section of follicles in the late active stage; C: Section of ripe oocytes in the ripe stage; D: Section of follicles in the partially spawned stage; E: Section of follicles in the recovery stage. Scale bars = 50 μ m. Abbreviations: LG, lipid granule; LM, lumen; MT, mesenchymal tissue; N, nucleus; NO, nucleolus; OC, oocyte; OG, oogonium; V, vacuole; YG, yolk granule.

TABLE 1. The shell height and first sexual maturity of female *Neptunea* (*Barbitonia*) *arthritis cumingii* from May to August, 2002.

Shell height (mm)	Gonadal developmental stage						Mature (%)
	EA	LA	RI	PS	RE	Total	
31.4–40.9	34					34	0.0
41.0–50.9	26	2	2			30	13.3
51.0–60.9	15	2	10	5		32	53.1
61.0–70.9		3	21	6		30	100.0
71.0–80.9		2	22	9		33	100.0
81.0–90.5			16	12		28	100.0
Total						187	100.0

Abbreviations: EA, early active stage; LA, late active stage; RI, ripe stage; PS, partially spawned stage; RE, recovery stage.

oocytes, as well as previtellogenic oocytes in the follicles (Fig. 7E). The individuals in this stage appeared from May to August, and the main spawning occurred between June and July when the seawater temperature rose to approximately 16–23°C.

Recovery Stage: After spawning, the undischarged vitellogenic oocytes in the lumen of the follicle undergo cytolysis, and each follicle was contracted, and then degeneration or resorption of undischarged vitellogenic or mature oocytes occurred. Thereafter, the rearrangement of newly formed connective tissues, a few oogonia and previtellogenic oocytes appeared on the newly formed follicular walls (Fig. 7F). The individuals in the recovery stage appeared from June to August.

First Sexual Maturity

Before and after spawning, a total of 187 female individuals (31.4–40.9 mm in shell height) were histologically examined to certify whether they reached maturity and participated in reproduction. The rate of shells of different sizes that reached first sexual maturity is summarized in Table 1. The breeding season of this species was from May to August. In the case of some individuals with gonad developmental stage in the late active stage in May through August, it is supposed that they reach maturity, except for individuals in the early active stage during the breeding season. First sexual maturity was 0% in female snails of 31.4–40.9 mm in shell height if they were at the early active stage during the breeding season.

The percentage of first sexual maturity of the female snail of 41.0 to 50.9 mm in shell height

was 13.3%. The percentages of first sexual maturity of the female individuals of 51.0 to 60.9 cm in shell height were over 50%, all of which were at the late active, ripe or partially spawned stages. First sexual maturity was 100% for snails over 61.0 mm in height.

DISCUSSION

Germ Cell Development and Vitellogenesis

As vitellogenesis commences, nuclei of the oocytes increased in size. Early vitellogenesis is characterized by proliferation of endoplasmic reticulum and mitochondria, both of which are closely associated with lipid droplets. According to our electron microscope observations of early vitellogenic oocytes of *N. arthritis cumingii*, the Golgi apparatus is thought to be involved in a number of vacuoles and small vesicles in the perinuclear region in the cytoplasm, with carbohydrate (glycogen) particles filling the vacuoles. Lipid droplets and lipid yolk granules are then added to the vacuoles and vesicles formed by the Golgi complex (referred as autotrophic by Taylor & Anderson, 1969), as in *Ilyanassa obsoleta* (Taylor & Anderson, 1969), *Biomphalaria glabrata* (deJong-Brink et al., 1976), *Mytilus edulis* (Reverberi, 1971), *Rapana venosa* (Chung et al., 2002), *Siphonaria capensis* (Pal & Hodgson, 2002), *Patella barbara*, *P. argenvillei*, *P. granularis*, *P. ocululus*, *P. miniata*, and *Helcion pectunculus* (Hodgson & Eckelbarger, 2000). This study suggests that the Golgi complex and various sizes of vacuoles are involved in the formation of lipid droplets in the early vitellogenic

oocytes. From our observations of oogenesis, it is assumed that the mitochondria and the endoplasmic reticulum near lipid droplets are involved in the formation of lipid droplets in the early vitellogenic oocyte. However, we did not find pinocytotic tubules, which are thought to be involved in yolk production as seen in the vitellogenic oocytes of *Agriolimax reticulatus* (Hill & Bowen, 1976; Dohmen, 1983). In the late vitellogenic oocyte, we also did not observe microvilli on the vitelline envelope, which is thought to be involved in helping in absorption, transportation and secretion of egg envelopes (Nørrevang, 1968), as seen in *Mactra chinensis* (Chung, 1997), *M. veneriformis* (Chung & Ryou, 2000), and *Siphonaria seriata* (Pal & Hodgson, 2002).

Formation of cortical granules is a prominent feature of late vitellogenic oocytes in most bivalves, such as *Mactra chinensis* (Chung, 1997) and *M. veneriformis* (Chung & Ryou, 2000). Regarding formation of cortical granules during oogenesis, Hodgson & Eckelbarger (2000) described that Golgi complexes appeared predominately in the cortical region of the ooplasm and secrete electrone-dense, cortical granule-like organelles in the vitellogenic oocytes of *Patella barbara*, and they stated that Golgi complexes synthesize cortical granules. In the present study, however, such structures were not observed in the vitellogenic oocytes, as in *Iliana obsoleta* (Taylor & Anderson, 1969) and *Rapana venosa* (Chung et al., 2002). Compared with *Patella barbara*, the lack of these structures is a prominent characteristic during oogenesis, representing a significant difference in *N. arthritica cumingii*. In the present study, proteid yolk granules, which appeared near the rough endoplasmic reticulum and modified mitochondrial structure (multivesicular bodies), as seen in *Hypselodoris tricolor* and *Godiva banyulensis* (Medina et al., 1986), were observed at the cortical region of the cytoplasm. Accord-

ingly, it is assumed that the endoplasmic reticulum and multivesicular bodies are involved in the formation of proteid yolk granules (Taylor & Anderson, 1969) as yolk precursor. In the present study, although the follicle cell, which lies adjacent vitellogenic oocyte, contains electron-dense granules and lipid droplets, we could not observe clear evidence of secretion into the vitellogenic oocyte. Therefore, it is assumed that *N. arthritica cumingii* synthesize yolk autotynthetically, as in the majority of gastropods, exceptions being some gastropods (*Planorbarius corneus*, *Lymnaea stagnalis*, *Hypselodoris tricolor*, *Godiva banyulensis*, *Siphonaria capensis*, and *S. serrata*) that synthesize yolk autotynthetically and heterotynthetically (Bottke et al., 1982; Medina et al., 1986; Pal & Hodgson, 2002).

Gonadal Development and Maturation

We observed that gametogenesis of *N. arthritica cumingii* begins at a temperature of about 3°C, with maximum gonadal maturation occurring in April 2002 and 2003, when water temperatures rose (Fig. 7) and phytoplankton was very abundant. Periods of high food abundance and gonad development were nearly coincident. In Korean coastal waters, growth and production of *Meretrix lusoria* and *Ruditapes philippinarum* are very high in the spring and early summer seasons (Kim et al., 1977; Chung et al., 1994; Lee, 1995) due to the abundant phytoplankton that occurs with increasing water temperatures. *Ruditapes philippinarum*, *Meretrix lusoria*, and other clams are commonly used as food by *N. arthritica cumingii*. At this time, abundant food can be supplied to *N. arthritica cumingii* during the period of gonadal development and maturation. Therefore, it is suggested that gonadal development and maturation of *N. arthritica cumingii* is closely related to water temperature and food availability.

TABLE 2. Comparisons of the spawning season of Buccinidae in each locality.

Species	Spawning season	Locality	Author
<i>Neptunea arthritica cumingii</i>	May–August	Kunsan, Korea	Present study
<i>N. cumingii</i>	July–August	East China Sea, China	Amio, 1963
<i>N. arthritica</i>	May–June	Usu Bay, Hokkaido, Japan	Fujinaga, 1985
<i>N. arthritica</i>	May–August	Saroma, Hokkaido, Japan	Kawai et al., 1994
<i>N. constricta</i>	December	East Sea, Korea	Son, 2003
<i>Siphonalia assidariaeformis</i>	December	East China Sea, China	Habe, 1960

Breeding Pattern

As shown in Table 2, our histological observations show that spawning of *N. arthritica cumingii* on the west coast of Korea occurs from late May to August in 2002 and 2003 when sea water temperatures were high. The spawning season of *N. cumingii* collected by the trawl net in the East China Sea occurs between July and August (Amio, 1963). *Neptunea arthritica* in Japan has been reported to spawn once a year between May and June in Utsu Bay, Japan (Fujinaga, 1985).

Therefore, it is assumed that the spawning period of *N. arthritica cumingii* on the west coast of Korea occurred somewhat earlier than that in the East China Sea. On the whole, *N. arthritica cumingii* in Korea is a summer breeder, based on the criteria outlined by Boolootian et al. (1962) for marine mollusks. In general, it is assumed that spawning of *N. arthritica cumingii* and *N. arthritica* in Korea and Japan occurs between May and August. However, spawning of *N. constricta* and *Siphonalia assidariaeformis* (Buccinidae) occurs during December, these species being winter breeders (Table 2). Therefore, the slight discrepancy in the spawning period between these studies might be related to geographic differences in water temperature and food availability (Chung et al., 2002).

First Sexual Maturity with the Gonad Developmental Stage

From the result of histological observations, we found that although the specimens were collected during the breeding season, the gonadal development of smaller individuals ranging from 31.4 to 40.9 mm in shell height were in the early active stage as small number of oogonia and the previtellogenic oocytes were present in the follicle of the ovary. Judging from histological observations, it is supposed that the size of the oocyte could not have reached maturity until late August, when spawning ended. Snails of 51.0–60.9 mm high were in the late active, ripe and partially spawned stages, and more than 50% reached first sexual maturity. However, all snails in the late active, ripe, or partially spawned stages spawned if they were larger 61.0 mm. This means that larger individuals can reach maturity earlier than smaller individuals. In the aspect of natural resources management, the present study suggests that because harvesting snails < 51.0 mm can potentially cause a

drastic reduction in recruitment, a prohibitory measure should be taken for adequate resource management. Henceforth, age determination by size of the individuals should be investigated in detail for natural resources management of this species.

ACKNOWLEDGEMENTS

The authors are grateful to Dr. John B. Burch of the University of Michigan for helpful comments on the manuscript. This research was supported in part by the fund (2003) from the Coastal Research Center, Kunsan National University.

LITERATURE CITED

- AMIO, M., 1963, A comparative embryology of marine gastropods, with ecological emphasis. *Journal of Shimonoseki College of Fisheries*, 12: 229–253.
- BOOLOOTIAN, R. A., A. FARMANFARMAINA & A. C. GIESE, 1962, On the reproductive cycle and breeding habits of two western species of *Haliotis*. *Biological Bulletin*, 122: 183–192.
- BOTTKE, W., I. SINHA & I. KEIL, 1982, Coated vesicle-mediated transport and deposition of vitellogenic ferritin in the rapid growth phase of snail oocytes. *Journal of Cell Science*, 53: 173–191.
- CHUNG, E. Y., 1997, Ultrastructural study of germ cell development and reproductive cycle of the hen clam, *Macra chinensis* on the west coast of Korea. *Development & Reproduction*, 1(2): 141–156.
- CHUNG, E. Y. & S. Y. KIM, 1996, Reproduction of *Neptunea arthritica cumingii* (Gastropoda: Buccinidae) on the west coast of Korea. *Proceedings of the Fourth International Congress on Medical and Applied Malacology*, 8(1): 170–171.
- CHUNG, E. Y., S. Y. KIM, K. H. PARK & G. M. PARK, 2002, Sexual maturation, spawning, and deposition of the egg capsules of the female purple shell, *Rapana venosa* (Gastropoda: Muricidae). *Malacologia*, 44: 241–257.
- CHUNG, E. Y. & D. K. RYOU, 2000, Gametogenesis and sexual maturation of the surf clam *Macra veneriformis* on the west coast of Korea. *Malacologia*, 42: 149–163.
- CHUNG, E. Y., D. K. RYOU & J. H. LEE, 1994, Gonadal development, age and growth of the shortnecked clam, *Ruditapes philippinarum* (Pelecypoda: Veneridae), on the coast of Kimje, Korea. *The Korean Journal of Malacology*, 10: 38–54.
- DE JONG-BRINK, M. A. WIT, G. KRAAL & H. H. BOER, 1976, A light and electron microscope study on oogenesis in the freshwater pulmonate snail *Biomphalaria glabrata*. *Cell and Tissue Research*, 171: 195–219.

- DOMEN, M. R., 1983, Gametogenesis. Pp. 1–48, in: N. H. VERDONK, J. A. M. VAN DEN BIGGELAAR & A. S. TOMPA, eds., *The Mollusca*, Vol. 3. Academic Press, New York.
- FUJINAGA, K., 1985, The reproductive ecology of the neptune whelk (*Neptunea arthritica* Bernardi) population, with special reference to the reproductive cycles, depositions of egg masses and hatchings of juveniles. *Bulletin of Faculty of Fisheries Hokkaido University*, 36: 87–98 [in Japanese with English summary].
- FUJINAGA, K., 1987, On the growth pattern of the neptune whelk, *Neptunea arthritica* Bernardi. *Bulletin of Faculty of Fisheries Hokkaido University*, 38: 191–202.
- GRIFFOND, B., 1980, La vitellogenèse chez la paludine *Viviparus viviparus* L. (Mollusque Gastéropode Prosobranchie). *Bulletin de la Société Zoologique de France*, 105: 401–407.
- GRIFFOND, B. & L. GOMOT, 1979, Ultrastructural study of the follicle cells in the freshwater gastropod *Viviparus viviparus* L. *Cell and Tissue Research*, 202: 25–32.
- HILL, R. S. & J. P. BOWEN, 1976, Studies on the ovotestis of the slug *Agriolimax reticulatus* (Müller). 1. The oocyte. *Cell and Tissue Research*, 173: 468–482.
- HODGSON, A. N. & K. J. ECKELBARGER, 2000, Ultrastructure of the ovary and oogenesis in six species of patellid limpets (Gastropoda: Patellogastropoda) from South Africa. *Invertebrate Biology*, 119: 265–277.
- ITO, H., 1982, Distribution of a sea snail, *Neptunea arthritica*, and its surroundings in the Lagoon Furen-ko. *Reports of the Engineering Research on all-round Constructions of Fishing Grounds in the Region of Nemuro Bay*: 107–114 [in Japanese].
- ITO, H. & S. TACHIZAWA, 1981, An estimation of the density of the available stock of a sea snail, *Neptunea arthritica*, by trap fishing. *Bulletin of Hokkaido Region Fisheries Research Laboratory*, 46: 113–119.
- KAWAI, K., S. YAMAGUCHI, N. IDE, S. GOSHIMA & S. NAKAO, 1994, Reproductive cycle and parasites infection in the neptune whelk *Neptunea arthritica* in Lagoon Saroma. *Venus*, 53: 105–112.
- KIM, B. A., B. K. KIM & S. U. KIM, 1977, Studies on the growth of hard clam, *Meretrix lusoria*. *Bulletin of Fisheries Research and Development Agency*, 17: 77–85.
- KWON, O. K., G. M. PARK & J. S. LEE, 1993, *Colored shells of Korea*. Academy Publishing Company, Seoul. 285 pp. [in Korean].
- LEE, Y. H., 1995, Growth and maturation of shortnecked clam, *Ruditapes philippinarum* by different seeding production areas. *Masters thesis, National Fisheries University of Pusan*. 51 pp. [in Korean].
- MACINTOSH, R. A. & A. J. PAUL, 1977, The relation of shell length to total weight, edible-meat weight, and reproductive organ weight of the gastropods *Neptunea heros*, *N. lyrata*, *N. prabilloensis*, and *N. ventricosa* of the eastern Bering Sea. *Proceedings of National Shellfish Association*, 67: 103–112.
- McCANN-COLLIER, M., 1977, An unusual cytoplasmic organelle in oocytes of *Ilyanassa obsoleta*. *Journal of Morphology*, 153: 119–127.
- McCANN-COLLIER, M., 1979, RNA synthesis during *Ilyanassa* oogenesis. An autoradiographic study. *Development, Growth and Differentiation*, 21: 391–399.
- MEDINA, A., J. C. GARCIA, F. J. MORENO & J. L. LOPEZ-CAMPOS, 1986, Comparative studies on the histology of the ovotestis in *Hypselodoris tricolor* and *Godiva banyulensis* (Gastropoda, Opisthobranchia), with special reference to yolk formation. *Journal of Morphology*, 188: 105–118.
- MIYAWAKI, M., 1953, Some observations on spawning of the whelk *Neptunea arthritica* Bernardi. *Zoological Magazine*, Tokyo, 62: 199–201 (in Japanese with English summary).
- NØRRVANG, A., 1968, Electron microscopic morphology of oogenesis. *International Review of Cytology*, 23: 113–186.
- PEARCE, J. & G. THORSON, 1967, The feeding and reproductive biology of the red whelk, *Neptunea antiqua* (L.). (Gastropoda, Prosobranchia). *Ophelia*, 4: 277–314.
- PAL, P. & A. N. HODGSON, 2002, An ultrastructural study of oogenesis in a planktonic and a direct-developing species of *Siphonaria* (Gastropoda: Pulmonata). *Journal of Molluscan Studies*, 68: 337–344.
- REVERBERI, G., 1971, *Mytilus* in experimental embryology of marine and freshwater invertebrates. Pp. 175–187, in: G. REVERBERI, ed., *Experimental embryology of Marine and freshwater invertebrates*. North-Holland, Amsterdam & London. xxiii + 587 pp.
- SON, M. H., 2003, Eggs and egg capsule morphology of the neptune whelk, *Neptunea constricta* (Dall, 1907) (Gastropoda: Buccinidae). *The Korean Journal of Malacology*, 19: 111–115.
- SUZUKI, K., T. HIRAISHI, K. YAMAMOTO & K. NASHIMOTO, 1996, Age determination and growth analysis based on size-frequency histograms of whelk *Neptunea arthritica* in Shiraiuchi, Hokkaido. *Nippon Suisan Gakkaishi*, 62: 225–229.
- TAKAHASHI, N., K. TAKANO & S. MURAI, 1972, Histological studies on the reproductive cycle of the male neptune whelk, *Neptunea arthritica*. *Bulletin of Faculty of Fisheries Hokkaido University*, 31: 314–326 [in Japanese with English summary].
- TAKAMARU, N. & A. FUJI, 1981, Reproductive cycle of the neptune whelk, *Neptunea arthritica* (Bernardi), in southern Hokkaido. *The Aquiculture*, 29: 78–87 [in Japanese with English summary].
- TAYLOR, G. T. & E. ANDERSON, 1969, Cytochemical and fine structural analysis of oogenesis in the gastropod, *Ilyanassa obsoleta*. *Journal of Morphology*, 129: 211–248.
- YOO, J. S., 1976, *Korean shells in color*. Ilgisa, Seoul. 196 pp. [in Korean].